

**KILLS
JAPANESE
BEETLES**



GARDEN TECH
Sevin® 5
READY-TO-USE
5% DUST



KILLS OVER 65 INSECTS
**ON VEGETABLES, FRUITS,
ORNAMENTALS & LAWN**

**STOP -
KEEP OUT OF REACH OF CHILDREN
CAUTION**

ACTIVE INGREDIENT:
Carbaryl (1-naphthyl N-methylcarbamate) 5.0%
OTHER INGREDIENTS: 95.0%
TOTAL: 100.0%

See Back Panel For Precautionary Statements
**PRECAUCION AL USUARIO: Si usted no puede leer o
entender inglés, no use este producto hasta que la
etiqueta le haya sido explicada ampliamente.**

Net Weight: 1 lb. (454 g)

Distributed by: TechPac, LLC,
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

Sevin is a registered trademark of
Tessenderlo Kerley, Inc.
Gardentech is a registered trademark
of Gulfstream Home and Garden Inc.

EPA Reg. No. 432-1209-71004

EPA Est. 67572-GA-1

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

68703-GA-1

68703-GA-2

FOR OUTDOOR RESIDENTIAL USE ONLY

FIRST AID

This product is a N-methyl carbamate and is a cholinesterase inhibitor

If swallowed:

- Call poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

If on skin or clothing:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-420-9347 for emergency treatment information.

NOTE TO PHYSICIAN: CARBARYL is an N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION
Harmful if swallowed. Harmful if absorbed through skin. Avoid contact with eyes, skin, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Remove contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS
This pesticide is extremely toxic to aquatic invertebrates. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

BEE CAUTION: MAY KILL HONEY BEES AND OTHER BEES. This product is highly toxic to honey bees and other bees. Do not apply to plants in bloom.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Wearing household latex or rubber gloves, use this ready-to-use shaker container to apply product. Dust lightly to cover leaf surfaces with a thin, even film of dust. Apply when air is calm. Start applying at the farthest corner of the treatment area and work backward. For plants above chest height, consider using a liquid carbaryl product.

RESTRICTIONS
Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

Non-refillable container. Do not cut package to expose contents.
Discard package when empty.

Not for use on plants grown for sale or commercial use, research, or commercial seed production.

PRECAUTIONS
Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days duration following application. Begin applications when insect or damage is first observed.

HOME GARDEN VEGETABLES AND SMALL FRUITS
Do not repeat applications more than once a week (7 days).

PREHARVEST USE RESTRICTIONS:
Observe the following preharvest intervals (number of days between last application and harvest)

PREHARVEST INTERVAL	
GARDEN CROP	PREHARVEST INTERVAL (DAYS)
Garden beets (roots), radishes, rutabagas, turnips (roots), carrots, potato, blackberries, boysenberries, dewberries, loganberries, raspberries, blueberries, grapes, and strawberries	7
Collards, garden beets (tops), kale, lettuce (head and leaf), mustard greens, spinach and turnip (tops)	14
Dried shelled legume beans and peas including: field bean, kidney bean, dry lima bean, navy bean, pinto bean, black-eyed pea, cowpea, Crowder pea, mung bean, Southern pea, lentil (dry).	21
Edible-podded legume vegetables including: Snap bean, wax bean, yardlong bean, jack bean, edible-pod pea, snow pea, sugar snap pea.	
Dried shelled legume vegetables including: field bean, kidney bean, lima bean (dry), navy bean, pinto bean, adzuki bean, black-eyed pea, cowpea, mung bean, Southern pea, lentil (dry). (Do not use on lentils in CA.)*	
Edible-podded legume vegetables including: Snap bean, wax bean, yardlong bean, jack bean, edible-pod pea, snow pea, sugar snap pea, broccoli, brussel sprouts, cabbage, cauliflower, kohlrabi, cantaloupe, cucumber, melons, pumpkins, squash, eggplant, pepper, and tomatoes	3

CROP	VEGETABLES	
	INSECT PESTS	COMMENTS
Edible-podded legume vegetables including: Snap bean, wax bean, yardlong bean, jack bean, edible-pod pea, snow pea, sugar snap pea.	Bean leaf beetle, cucumber beetles, corn earworm, flea beetles, Japanese beetles, leafhoppers, Lima bean pod borer, Lygus bug, Mexican bean beetle, stink bugs	Maximum number of applications per year: 4
Broccoli, brussel sprouts, cabbage, cauliflower, kohlrabi	Amyworms, corn earworm, diamond-back moth larvae, Flea beetles, harlequin bugs, imported cabbageworm	Application is only permitted within thirty days from the date of crop emergence or the date of transplanting. Will not control cabbage loopers. Maximum number of applications per year: 4

CROP	VEGETABLES - CONTINUED	
	INSECT PESTS	COMMENTS
Collards, garden beets, kale, lettuce, mustard greens, radishes, rutabagas, spinach, turnips	Amyworms, corn earworm, Diamond-back moth larvae, Flea beetles, harlequin bug, imported cabbageworm, leafhoppers, stink bugs, tarnished plant bug	Application is only permitted within thirty days from the date of crop emergence or the date of transplanting. Will not control cabbage loopers. Maximum number of applications per year: 4
Cantaloupe, cucumbers, melons, pumpkins, squash	Cucumber beetles, flea beetles, melonworms, pickleworm, squash bug	Maximum number of applications per year: 6
Carrots	Flea beetles, leafhoppers, lygus bugs, spittlebugs, stink bugs	Maximum number of applications per year: 6
Eggplants, pepper, tomato	European corn borer, Fall armyworm, flea beetles, leafhoppers, tomato fruitworm, tomato hornworms, stink bugs	Maximum number of applications per year: 7
Potato	Colorado potato beetle, flea beetles, leafhoppers, lygus bugs	Maximum number of applications per year: 6

***Not for use on fresh/succulent peas and beans.**

PLANT	SMALL FRUIT	
	INSECT PESTS	COMMENTS
Blackberries, boysenberries, strawberries, loganberries, dewberries, raspberries	Japanese beetles, leafhoppers, leafrollers, raspberry aphid	Maximum number of applications per year: 5
Blueberries	Blueberry maggot, cherry fruitworms, cranberry fruitworms, Japanese beetles	Apply 3 weeks before harvest and repeat 10 days later if necessary. Maximum number of applications per year: 5
Grapes	Grape berry moths, grape leafroller, grape leafhopper, grape leaf skeletonizer, Japanese beetle, June beetle, red-banded leafroller	Apply just before first brood leafroller larvae emerge from rolls and as needed for leafhoppers. Maximum number of applications per year: 5
Strawberries	Meadow spittlebug, strawberry leafroller, strawberry weevil	Maximum number of applications per year: 5

ORNAMENTAL SHRUBS AND FLOWERS
For outdoor use only. To kill bagworm, blister beetles, boxelder bug, boxwood leafminer, flea beetles, Japanese beetle, lacebugs, leafhoppers, leafrollers, periodical cicada, plant bugs, psyllids, rose aphid, rose slug, scale insects in the crawler stage, tent caterpillars and exposed thrips, apply thoroughly to the infested plants at first sign of damage. Do not repeat application more than once weekly or more than six times per year. For best results on scale, apply in spring and early summer when scale crawlers are present.

HOME LAWN
To kill earwigs, fall armyworm, leafhoppers, millipedes, sod webworm, and sowbugs, sprinkle lightly on grass.

To kill ants (except pharaoh and carpenter ants), chiggers, fleas, spiders (except Brown recluse and Black widow) and ticks, sprinkle lightly on grass. Will only kill pests present at the time of application. Pest must be directly contacted by product to be effective.

For best results, apply after rain or watering and do not water for at least 2 days after application. Repeat treatment as needed but not more than every 2 weeks and not more than four times per year.

EARWIGS
This material is effective in killing earwigs and when properly used will reduce the number gaining entrance to homes. Using this ready-to-use shaker container, lightly dust a band 3 to 4 inches wide around the outside foundation wall of the home. Earwigs breed over wide areas around yards and fields and are driven by cold weather to seek shelter in warm buildings. As they move along the foundation wall seeking entrance, they come into contact with the treated barrier and are killed. Be sure to treat in front of all steps and other entrance ways as well as along the foundation wall. Apply also around refuse piles, lumber, mulch and other areas in the yard offering areas of concealment to earwigs. Repeat application in 10 days or when earwigs become troublesome. If the dust barrier is washed away by rain, it should be replaced with a new application if earwigs are still a problem.

This container is sold by weight, NOT by volume. You can be assured of the proper weight even though some settling of contents normally occurs during transportation and handling.

Important: Read the entire Directions before using this product. Disclaimer: Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions for Use. To the extent consistent with applicable law, manufacturer makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. To the extent consistent with applicable law, in no case shall Manufacturer be liable for consequential, special or indirect damages resulting from the use, handling or application of this product.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage and disposal.
Pesticide Storage: Store pesticide in original container and place in a locked storage area.
Pesticide Disposal: If empty: Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. If Partly Filled: Call your local solid waste agency or 1-866-945-5033 for disposal instructions. Never place unused product down any indoor or outdoor drain.

NOTICE: Cancer - www.P65Warnings.ca.gov

