



PHOSPHORUS-FREE

PHOSPHORUS-FREE

PHOSPHORUS-FREE

# WEED & FEED

Use on lawns to kill the toughest weeds, like clovers and dandelion

- ◆ Controls a wide spectrum of broadleaf weeds
- ◆ Quick greening and extended feeding

FEEDS UP TO 3 MONTHS

**WEED & FEED**  
Kills the Toughest Weeds

**WEED & FEED**  
Kills the Toughest Weeds



UPC

UPC

<b>ACTIVE INGREDIENTS:</b>	
2,4-D, 2-ethylhexyl ester.....	1.108%
Mecoprop-p acid.....	0.167%
Dicamba acid.....	0.071%
<b>OTHER INGREDIENTS:</b> .....	98.654%
<b>TOTAL</b> .....	100.000%

**THIS PRODUCT CONTAINS:**  
 14.70 lb. = 2,4-dichlorophenoxyacetic acid equivalent per ton or 0.735%, CAS 1928-43-4.  
 3.34 lb. = (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton or 0.167%,  
 CAS 18484-77-8.  
 1.42 lb. = 3,6-dichloro-*o*-anisic acid equivalent per ton or 0.071%, CAS 1918-00-9.  
 Isomer Specific by AOAC Methods

KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See Back Panel For First Aid And Additional Precautionary Statements

**25-0-4**

SEE BACK PANEL FOR ANALYSIS DESCRIPTION

**NET WT 39 LBS (17.69 kg)**

**COVERS  
15,000 SQ FT**



PHOSPHORUS-FREE

# WEED & FEED

## GUARANTEED ANALYSIS

25-0-4

Total Nitrogen (N).....	25.0%
25% Urea Nitrogen*	
Soluble Potash (K <sub>2</sub> O).....	4.0%

Derived from: Urea, Polymer-Coated Urea, Muriate of Potash.

\*5.0% Slow Release Urea Nitrogen From Polymer-Coated Urea.

F525

**STOP** READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

**CAUTION:** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

### FIRST AID

**If in Eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a Poison Control Center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For information on this product, contact the National Pesticide Information Center, 1-800-858-7378, Monday-Friday, 8:00 a.m. to 12:00 p.m. PST. You may also contact the National Poison Control Center, 1-800-222-1222, day or night, for emergency medical treatment information. FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident call CHEMTREC 1-800-424-9300.

### Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

### PRECAUTIONS AND RESTRICTIONS:

- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.
- Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

**INSTRUCTIONS:** This product is a combination of three proven weed killers -- 2,4-D, mecoprop-p, and dicamba. This herbicide controls a wide range of lawn weeds, including Dandelion, Chickweed, Knotweed, Plantain, Henbit and Spurge. In addition, a 25-0-4 fertilizer provides a supply of plant nutrients.

### USE PRECAUTIONS

Do not use on dichondra, carpetgrass, "Floramat" St. Augustinegrass, creeping bentgrass, or turfgrass where desirable clovers are present.

Do not use this product to newly seeded grasses until well-established or after the second or third mowing.

Do not use this product on flowers, vegetables, groundcovers, gardens or landscape plantings including shrubs and trees. Do not exceed specified dosage for any area and do not apply this product within the drip line of, or over surface roots of trees, shrubs, and other ornamentals.

Re-seeding or overseeding areas may occur three (3) weeks after the application of this product.

Do not use this product on or near desirable plants and avoid spreading granules onto exposed root systems or adventitious shoots (rhizomes) within the drip line of the roots of desirable trees and shrubs, since injury may result.

Do not apply directly to or near water, storm drains, gutters, sewers or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. Apply this product directly to your lawn, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

### WHEN TO APPLY

Apply when broadleaf weeds are actively growing in Spring or early Fall.

Apply when the air is calm to ensure uniform coverage and avoid spreading granules onto flowers, vegetables, and ornamental shrubbery.

Possible retreatment may be necessary if lawn is heavily overrun by weeds, if adverse low moisture conditions prevail, or if weeds are in a state of poor growth.

### HOW MUCH TO APPLY

For use on Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, common Bermudagrass, Bahiagrass, Zoysiagrass, St. Augustinegrass, Centipedegrass, hybrid Bermudagrass, and Colonial Bentgrass. Apply 2.6 pounds of product per 1,000 square feet of turfgrass. A second application may be made in approximately 30 days. The maximum number of broadcast applications per treatment site is 2 per year.

### APPLICATION INSTRUCTIONS

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 to 2 days before application to sustain moisture until the next watering (see step 5).
3. Apply when Broadleaf weeds are young and actively growing, preferably in the morning when dew is on the grass. At the time of application, moisture on the weed leaves from dew, rainfall or irrigation/watering aids in control. If grass is not moist at time of application, sprinkle lightly with water before applying product.
4. Avoid applying when temperatures are consistently above 90°F.
5. Do not wash from weed leaves for 1 to 2 days after application. At this time, a thorough watering should be made.

EPA Reg. No. 2217-884-59144  
EPA Est. No. 59144-GA-1(A), 32802-MO-1(B), 37351-CO-1(C), 8378-IN-1(D), 9404-FL-1(E), 961-PA-01(F), 961-IL-01(G), 35497-OR-1(H), 49991-TX-2(Q), 69759-OR-002(R), 7401-TX-01(S)  
Letter in parenthesis is first letter of lot number.

## HOW TO APPLY--SPREADERS AND CALIBRATION

Always apply with a rotary or drop spreader to distribute the granules evenly. Never apply by hand.

**COVERAGE** – To determine how much you need to apply, multiply the length of your lawn by its width. This equals your total square footage. Deduct all non-lawn areas (house, gardens, etc.). One bag covers up to 5000 square feet.

**Important:** It is important to follow directions as indicated on this quality lawn care product as over-application may cause injury to your lawn.

Spreader settings are approximate. Differences may occur due to condition of spreader, speed at which applied and roughness of area treated. Be sure to check spreader before beginning, adjusting if necessary. Consult Owner's Manual. Never apply by hand.

### SINGLE PASS DISTRIBUTION:

- For rotary spreaders: Operate at intervals corresponding to the swath width.
- For drop spreaders: Overlap the wheel tracks to ensure complete coverage.
- For best results, apply this product uniformly and avoid streaks, skips or overlaps. Walk at a steady pace to ensure even coverage. Close the spreader opening when turning to change directions or when stopping.

### CALIBRATION INSTRUCTIONS AND SPREADER SETTINGS:

Calibrate the spreader according to the manufacturer's directions. Initial spreader settings may require adjustment to deliver the recommended rate under actual application conditions.

To provide proper distribution, calibrate spreader before application.

• For 2.6 pounds of product per 1,000 square feet of turfgrass: Check your spreader on 385 square feet and adjust to apply at the rate of 1 pound of product per 385 square feet.

**BROADCAST SPREADERS:** For best results, apply in a circular path starting on the outside and working toward the center overlapping slightly to insure complete coverage.

**DROP SPREADERS:** Apply 2 strips along each end. Then apply along length of lawn, overlapping slightly and shutting off the spreader before each turn at edge of border strips.

SPREADER SETTINGS			
BROADCAST SPREADER		DROP SPREADER	
Scotts SpeedyGreen	4.5	Scotts AccuGreen	6
Scotts Basic	4.5		
Scotts Lawn Pro	4.5		
Scotts EdgeGuard Deluxe	4.5		
Scotts EdgeGuard Mini	4.5		
Precision Rotary	5		
EV-N-SPRED	13		

### This product controls the following list of growing weeds:

Amaranth, purple	Chickweed, mouseear	Fleabane, rough	Mustard, hedge	Ragweed, bur	Thistle, yellow (horrible)
Amaranth, slender	Chickweed, sticky	Galinsoga, hairy	Mustard, indian	Ragweed, common	Tailing crownvetch
Aster, heath	Chicory	Galinsoga, smallflower	Mustard, tansy	Ragweed, lanceleaf	Velvetleaf
Aster, white heath	Cinquefoil	Goldenrod	Mustard, tumble	Ragweed, western	Vervain, blue
Aster, purple	Clover, crimson	Ground ivy	Mustard, white	Redstem filaree	Vervain, hoary
Aster, white prairie	Clover, alyce	Groundsel	Mustard, wormseed	Rough cinquefoil	Vervain, prostrate
Artichoke	Clover, hop	Gumweed	Nettle, stinging (burning)	Scarlet pimpernel	Vervain, tall
Austrian fieldcress	Clover, rabbit foot	Hairy fleabane	Nettle, tall	Shepardspurge	Vetch, bird
Bedstraw, Catchweed	Clover, red	Hawkweed, meadow	Nettle, wood	Smartweed, ladythumb	Vetch, hairy
Bedstraw, smooth	Clover, strawberry	Hawkweed, mouseear	Parsley-piert	Smartweed, pale	Vetch, milk
Bedstraw, Northern	Clover, yellow sweet	Hawkweed, orange	Parson	Smartweed, pennsylvania	Vetch, narrowleaf
Bedstraw, pineywoods	Clover, white sweet	Hawkweed, yellow	Pearwort	Smooth chaff-flower	Vetch, wild
Beggarticks, hairy	Clover, white	Healall	Pennycress (frenchweed)	Smoothdock	Virginia creeper
Betony, Florida	Cockle	Heartleaf drymary	Pepperweed, perennial	Sorel, red (sheep)	Western clematis
Bindweed, Field	Cocklebur	Hemp	Pepperweed, virginia	Sowthistle, annual	Western salsify
Bindweed, Hedge	Cocklebur, oriental	Henbit	(Peppergrass)	Sowthistle, spiny	Wild aster
Bitter Wintercress	Cocklebur, spiny	Hoary cress	Pigweed, prostrate	Spanish needles	Wild buckwheat
Bittercress, hairy	Common mullein	Horse-nettle	Pigweed, redroot	Speedwell, bi-lobed	Wild carrot
Bitterweed	Corn Spurry	Horseweed	Pigweed, russian	Speedwell, common	Wild four-o'clock
Black-eyed Susan	Creeping Beggarweed	Jimsonweed	Pigweed, smooth	Speedwell, corn	Wild garlic
Black Medic	Creeping Jenny	Johnny-jumpup violet	Pigweed, tumble	Speedwell, ivyleaf	Wild lettuce
Blood flower milkweed	(creeping charlie)	Knawel	Pineapple weed	Speedwell, persian	Wild marigold
Blue lettuce	Cupid shaving Brush	Knotweed, prostrate	Plains coreopsis (tickseed)	Speedwell, purslane	Wild mustard
Brassbuttons	Cudweed	Kochia	Plantain, blackseed	Speedwell, slender	(yellow rocket)
Bristly oxtonogwe	Daisy, English	Lambsquarters	Plantain, bracted	Speedwell, snow	Wild onion
Broomweed	Daisy, oxeye	Lespedeza	Plantain, broadleaf	Speedwell, thymeleaf	Wild parsnip
Buckhorn	Dandelion, common	Mallow, alkali	Plantain, buckhorn	Spiny amaranth	Wild radish
Bullnettle	Dandelion, false	Mallow, bristly	Plantain, hoary	Spurge, spotted	Wild rape
Burclover	Day flower	Mallow, common	Plantain, narrowleaf	Spurge, prostrate	Wild strawberry
Burdock	Deadnettle	Mallow, dwarf	Plantain, slender	Spurweed	Wild sweet potato
Burweed, lawn	Dichondra	Mallow, venice	Plantain, woolly	Star of Bethlehem	Wild violet
(spurweed)	Dock, broadleaf	Marcela	Poison ivy	Strawberry, India mock	Woodsorel, creeping
Buttercup, bulbous	Dock, curly	Matchweed	Poison oak	Tansy ragwort	Woodsorel, violet
Buttercup, creeping	Doggbane	Mexicanweed	Pokeweed	Tanweed	Woodsorel, yellow (Xnals)
Butterweed, common	Dogfennel	Moneywort	Prairie Sunflower	Thistle, blessed	Woolly croton
(poorjoe)	Dollarweed (Pennywort)	Morningglory, bigroot	Prickly lettuce	Thistle, bull	Wormseed
Buttonweed, virginia	Elderberry	Morningglory, ivyleaf	(compass plant)	Thistle, canada	Yarrow, common
Carolina geranium (wild)	Falseflax	Morningglory, mexican	Prickly sida	Thistle, fiodman (prairie)	Yarrow, western
Carpetweed	Filaree, redstem	Morningglory, red	Prostrate knotweed	Thistle, musk	Yarrow, yellow
Catnip	False sunflower	Morningglory, tall	Puncturevine	Thistle, platte	
Catsear, spotted	Fiddleneck	Morningglory, wolly	Purslane, common	Thistle, plumeless	
(false dandelion)	Filaree, redstem	Mugwort Mustard, ball	Pusley, brazil	Thistle, russian	
Chamber bitter	Filaree, whitestem	Mustard, black	Pusley, florida	Thistle, scotch	
Chickweed, common	Fleabane, daisy (annual)	Mustard, blue	Pusley, large flower	Thistle, wayleaf	

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. If Empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Keep this product off sidewalks, patios, driveways, etc. This fertilizer may stain concrete and painted surfaces. If it comes in contact with concrete surfaces, sweep off immediately before flushing with water. If concrete is wet when contact occurs, staining may be impossible to prevent. Be careful when using rotary spreaders which scatter fertilizer beyond lawn edge and onto concrete.

When applied as directed, this product meets the guidelines for metals adopted by the Association of American Plant Food Control Officials.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfc.org/metals.htm>.

### LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

Distributed & Guaranteed by  
Gro Tec, Inc. • P. O. Box 290 • Madison, GA 30650

U P C

Net Wt 39 lb (17.69 kg)